Table 5
Tule Marsh Habitat Vegetation Monitoring Results

raie maisii riabitat vegetation monitoring liesuits												
Monitored Element	Final Performance Standard	1st Year	2 nd Year	3 rd Year	4 th Year	5 th Year	6 th Year	7 th Year	8 th Year	9 th Year	10 th Year	
Extent	32 acres	3 acres	4 acres	7 acres	32 acres	34 acres	32 acres	33 acres	33.2 acres	>33.2 acres	>33.2 acres	
Absolute Cover	60%	0.02%	27%	100%	22.5%	40%	60%	65%	75%	±75%	±75%	
Relative Tall Emergent Cover	50%	N/A	N/A	N/A	80%	85%	50%	73%	75%	±75%	±75%	
Relative Native Cover	80%	100%	100%	81%	90%	95%	80%	100%	99.7%	±99%	±80%	
Plant Height	4 feet	N/A	N/A	N/A	2.5 ft.	4.0 ft.	4.0 ft.	7.8 ft.	8.7 ft.	±8.7 ft.	±8.7 ft.	
Relative Hydrophyte Cover	90%	100%	100%	100%	100%	99%	90%	100%	100%	100%	100%	
Open Water	<20%	N/A	N/A	N/A	>70%	<45%	<20%	<20%	<20%	<20%	<20%	



Photo 5: View from the northwest corner of the property looking south. June 2018.




Photo 6: View looking east from the northwest corner of the property. June 2018.

Based on aerial photographs, the extent of the tule marsh habitat has not changed since last year's results and remains above the performance standard of 32 acres. Since planting in 2012, the tules have exhibited steady, moderate growth. However, over the past two years the rate of growth appears to have slowed. This is likely due to the relatively high tule cover on those lands that were readily available and suitable.

Absolute cover within the tule marsh habitat remains high and above the performance standard of 60%. Based on a visual comparison of aerial photographs from last year and this year, absolute cover within the tule marsh habitat has remained approximately the same (between 70% and 80%). As discussed in previous years monitoring reports, the areas previously planted with tules have coalesced and now have almost continuous cover. These planted areas appear to be slowly expanding into the areas not previously planted, thus resulting in slow but consistent increases in absolute cover.

Relative tall emergent cover is high and above the performance standard of 50%. Although cover from primrose and other species is present throughout the sample plots, tules and a small number of cattails dominate the marsh. Plant height throughout the marsh has increased annually since planting and is significantly above the performance standard of 4 feet. Relative hydrophyte cover is very high at 100% and open water is less than 20%.

Invasive weed growth in this zone is seasonal and controlled by tides and wind. Generally, water hyacinth (*Eichorina crasspies*), six petal primrose (*Ludwigia hexapetala*) and other non-native aquatics reach peak growth in August to September and cover up to 10 to 15% of the total water area. This year, six petal primrose has been growing within the southern part of the site and cover increased from approximately 15% of the marsh at the time of monitoring in early summer to approximately 25% in early fall. Water hyacinth has, however, been close to absent from the site with less than 1% cover.

Shin Kee generally does not experience the explosive growth in aquatic weeds that other areas of the delta experience. This is generally because the marsh is oriented in a northwest-southeast direction, the same direction as the prevailing winds, which keeps the floating aquatics penned into the southeast edges of the site. With the advent of winter, these then die off and are flushed out through the culverts in the southeast corner or are stranded above the high tide level by the winter storms, where they then become part of the organic detritus in this zone. We expect that the six petal primrose growing within the southern part of the site will flush out though the culverts or get stranded above the high tide line this winter. We anticipate a substantial decline in six petal primrose cover next year and do not anticipate that the species will be as abundant in the coming years.

Although the tule marsh continues to meet its performance standards, we noticed a decline in tule cover in the southwestern corner of the marsh three years ago. We have been carefully monitoring this area and have noticed a slow return of tules to the area. To further document and monitor the return of tules to the area we placed six sample plots in this area and will monitor these plots for five years or until the tule population has returned and stabilized.

C. Photo Documentation

Photographs were taken from established photo points to document site conditions and are supplied throughout the text.

III. MANAGEMENT RECCOMENDATIONS

Monitoring and management over the past several years has focused primarily on the performance and extent of the tule marsh habitat; however, monitoring throughout the site has demonstrated the ability of the native plants to spread and out compete non-native competitors. Given the relatively stable performance of the habitats at Shin Kee over the past several years, this report recommends that maintenance continue to follow the current management plan including:

- 1. Mow levee every other year to reduce non-native vegetation. The levee was mowed during the summer of 2018 and should be mowed again in 2020.
- 2. Survey vegetation annually for exotic invasive species.
- 3. Survey six petal primrose within the southern part of the marsh to see if it is leaving the site through the southern culverts.
- 4. As needed, spot spray patches and hand pull exotic invasive vegetation to control population and prevent spread.

The hydrologic regime, as noted above, has produced optimal conditions for GGS and tule marsh over the past several years and will be left as is. The gates are now set at specific conditions that create average water surface elevations and tidal ranges similar to those found at the Coldani Marsh (which also has tide gates) and can likely be left at their current positions in perpetuity.

I. REPORT DISTRIBUTION LIST

The Westlake Villages Mitigation Monitoring Report is distributed to the following:

William Guthrie & Krystal Bell David Smith

Regulatory Branch EPA - Region 9

U.S. Army Corps of Engineers Attention: CWA Compliance

1325 J Street 75 Hawthorne Street

Sacramento, CA 95814 San Francisco, CA 94105-3901

Brian Hansen Andrea Boertien

USFWS Department of Fish and Game

Bay-Delta Fish & Wildlife Office Environmental Scientist

650 Capitol Mall, 8th Floor Bay-Delta Region Room 8-300

Sacramento, CA 95814 4001 N. Wilson Way

Stockton, CA 95205

Steve Mayo Peter Minkel

San Joaquin County Council of Environmental Scientist

Governments Regional Water Quality Control Board

555 East Weber Ave 11020 Sun Center Drive #200

Stockton, CA 95202 Rancho Cordova, CA 95670-6114

David Nelson Sean Skaggs

A. G. Spanos Companies Ebbin Moser + Skaggs LLP

10100 Trinity Parkway, 5th Floor 525 B Street, Suite 1500

Stockton, CA 95219 San Diego, CA 92101

Mr. David Smith, Wetlands Section Chief

APPENDIX A

Vegetation Monitoring Results

		Hydr	ophyte					S	hin Kee I	ntertidal	Marsh		
]	Nativ	e	Plot:	3	7	11	15	20	26	33	Ave.
Common Name	ROIND	<u> </u>		Per	Unveg:								0.00
yerba mansa slender aster	OBL FACW	1 1	$\frac{1}{1}$	1	Anemopsis californica Aster subulatus var. ligulatus					0.1			0.00
Santa Barbara sedge	FACW	1	1	1	Carex barbarae					0.1			0.01
clustered field sedge	FACW-	1	+i	1	Carex praegracilis					2.5			0.36
brown dogwood	FACW	1	1	1	Cornus glabrata								0.00
tall flat-sedge	FACW	1	1	1	Cyperus eragrostis								0.00
Sedge sp.	FACW	1	1	1	Cyperus sp.		0.1						0.01
salt grass spikeweed	FACW OBL	1	1	1	Distichlis spicata Eleocharis sp.				87.5				0.00 12.50
common waterweed	OBL	$\frac{1}{1}$	$+\frac{1}{1}$	 	Elodea canadensis				67.3				0.00
hairy willowherb	FACW	1	1	1	Epilobium ciliatum								0.00
seaside heliotrope	OBL	1	1	1	Heliotropium curassavicum							`	0.00
water penny	OBL	1	1	1	Hydrocotyl sp.								0.00
Baltic rush	OBL	1	1	1	Juneus balticus		1.5	2.5		27.5	60.5	27.5	0.00
soft rush iris-leaved rush	OBL OBL	<u> </u>	1 1	1	Juncus effusus Juncus xiphioides	 	15	2.5		37.5	62.5	37.5	22.14 0.00
creeping wildrye	FAC+	1	1	1	Leymus triticoides					15	87.5		14.64
water primrose	OBL	1	1	1	Ludwigia peploides		87.5	37.5	15	37.5	07.3		25.36
swamp smartweed	OBL	1	1	1	Polygonum hydropiperoides								0.00
Fremont cottonwood	FACW	1	1	1	Populus fremontii								0.00
California rose	FAC+	1	1	1	Rosa californica					2.5			0.36
gray willow	OBL	1	1	1	Salix exigua					1			0.14
Gooding willow red willow	OBL FACW+	1	1	1	Salix goodingii Salix laevigata							15	0.00 2.14
arrovo willow	FACW	1	$\frac{1}{1}$	1	Salix lasiolepis							13	0.00
tule	OBL	1	1	1	Scirpus acutus		15					 	2.14
three square	OBL	1	1	1	Scirpus americanus	15			62.5		15	2.5	13.57
salt marsh bullrush	OBL	1	1	1	Scirpus robustus	87.5							12.50
southern cattail	OBL	1	1	1	Typha domingensis				15	~=-	37.5	37.5	12.86
broad-leaved cattail	OBL	1	1	1	Typha latifolia	2.5	15	87.5		87.5			27.50 0.00
stinging nettle fat hen	FACW FACW	<u> </u>		0	Urtica dioica ssp holosericea Atriplex triangularis								0.00
horseweed	FACW	1	$\frac{1}{1}$	0	Convza canadensis								0.00
toad rush	FACW+	1	Ιi	0	Juncus bufonius	 					<u> </u>		0.00
broad-leaved pepperweed	FACW	1	0	1	Lepidium latifolium								0.00
bird's-foot trefoil	FAC	1	0	1	Lotus corniculatus								0.00
dallisgrass	FAC	1	0	1	Paspalum dilatatum		0.1		15	2.5			2.51
Vasey grass	OBL	1	0	1	Paspalum urvillei						ļ	2.5	0.36
Kentucky bluegrass dooryard knotweed	FAC FAC	1	0	1	Poa pratensis Polygonum arenastrum								0.00
water beard grass	FACW	1	0	1	Polypogon viridis					0.1			0.00
green dock	FACW	1	0	1	Rumex conglomeratus					0.1			0.00
curly dock	FACW-	1	0	1	Rumex crispus	 							0.00
Fiddleleaf dock	FAC	1	0	1	Rumex pulcher					0.1		0.1	0.03
Himalaya blackberry	FACW	1	0	1	Rubus discolor								0.00
Pacific bent grass	FACW*	1	0	0	Agrostis avenacea	ļ				1	2.5	37.5	5.71
Slender centaury lamb's quarters	FACW FAC	1	0	0	Centaurium tenuiflorum Chenopodium album					1	0.1		0.16
poison hemlock	FACW	1	0	0	Conium maculatum								0.00
water hyacinth	OBL	1	0	0	Eichhornia crassipes								0.00
weedy cudweed	FACW-	1	0	0	Gnaphalium luteo-album					0.1			0.01
Seaside barley	FAC	1	0	0	Hordeum marinum								0.00
prickly lettuce	FAC	1	0	0	Lactuca serriola								0.00
bearded sprangletop	OBL	1	0	0	Leptochloa fiscicularis								0.00
Italian ryegrass hyssop loosestrife	FAC FACW	1	0	0	Lolium multiflorum Lythrum hyssopifolia								0.00
India sweet-clover	FAC	1	0	0	Melilotus indica		2.5			2.5			0.71
bristly ox-tongue	FAC	1	0	0	Picris echioides					0.1	0.1	2	0.31
annual blue-grass	FACW-	1	0	0	Poa annua					<u> </u>			0.00
rabbitfoot grass	FACW+	1	0	0	Polypogon monspeliensis	000000000000000000000000000000000000000			0.1	15		2.5	2.51
coyote bush	UPL	0	1	1	Baccharis pilularis							I	0.00
big leaf mistletoe	UPL UPL	0	1	1	Phoradenron macrophyllum	ļ				0.1			0.00
parched willowherb Bindweed	UPL	0	1 0	0	Epilobium brachycarpum Convolvulus arensis					0.1			0.01
Bermuda grass	FACU	0	0	1	Cynodon dactylon	2.5		 			 		0.00
wild oat	UPL	0	0	0	Avena fatua		000000000000000000000000000000000000000	000000000000000000000000000000000000000	000000000000000000000000000000000000000				0.00
ripgut grass	UPL	0	0	0	Bromus diandrus								0.00
soft chess	FACU-	0	0	0	Bromus hordeaceus								0.00
Spanish Brome	UPL	0	0	0	Bromus madritensis								0.00
Italian thistle Big heron bill	UPL FACU	0	0	0	Carduus pycnocephalus	 	ļ	 	ļ		 	 	0.00
Rattail sixweeks grass	UPL	0	0	0	Erodium botrys Festuca myuros	 						 	0.00
Yellow sneezeweed	FACU	0	0	0	Helenium amarum							0.1	0.00
summer mustard	UPL	0	0	0	Hirschfeldia incana	<u> </u>	<u> </u>			<u></u>	<u></u>		0.00
Foxtail barley	FACU	0	0	0	Hordeum murinum								0.00
mallow	UPL	0	0	0	Malva sp								0.00
pineapple weed	FACU	0	0	0	Matricaria matricarioides								0.00
Bur clover	FACU+	0	0	0	Medicago polymorpha	-		-			 	 	0.00
white sweet-clover Wild radish	UPL	0	0	0	Melilotus alba Raphanus sativus	 						├──	0.00
common groundsel	NI	0	0	0	Senecio vulgaris	 		 			 		0.00
common sow-thistle	NI	0	0	0	Sonchus oleraceus						<u></u>		0.00
		**************	***************************************	<u></u>	(Sp. Rich.) Number of species:	4	7	3	6	17	7	10	7.71
	mid-point	s			Number of native species:	3	5	3	4	8	4	4	4.43
5=75-100%	5==87.5				Total veg cover	107.5	135.2	127.5	195.1	205.1	205.2	137.2	158.97
4=50-75%	3==62.5			<u> </u>	elative veg cover (=100-unveg):	 		 	ļ				
						100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.00
3=25-50%	3==37.5			<u> </u>	\(\sum_{\text{cover of all hydrophytic spp:}}\)	105.0	135.2	127.5	195.1	205.0	205.2	137.1	158.59
2=5-25%	2==15			<u> </u>	\sum cover of all native spp:	105.00	132.6	127.5	180.0	183.7	202.5	92.5	146.26
1=many-5%	1==2.5				\sum cover of all perennial spp:	107.5	132.7	127.5	195.0	186.2	202.5	95.0	149.49
d=few	d==0.1				relative cover of hydrophytes:	98%	100%	100%	100%	100%	100%	100%	100%
r=one	r==0.01				relative cover of natives:	98%	98%	100%	92%	90%	99%	67%	92%
				_									

		Hydro		<u> </u>	1	Shin Kee Riparian Wetland								
		11,5	Nativ	⁷ e	Plot:	4	8	10	17	22	27	34	Ave.	
Common Name	ROIND		1	Per	Unveg:		15					0.1	2.16	
yerba mansa slender aster	OBL FACW	1	1 1	1 1	Anemopsis californica Aster subulatus var. ligulatus			0.1	0.1				0.00	
Santa Barbara sedge	FACW	1	1	1	Carex barbarae	37.5	15	0.1	37.5	15	87.5	15	29.64	
clustered field sedge	FACW-	1	1	1	Carex praegracilis		37.5	87.5	37.5	37.5			28.57	
brown dogwood tall flat-sedge	FACW FACW	1	1	1	Cornus glabrata Cyperus eragrostis								0.00	
Sedge sp.	FACW	1	$\frac{1}{1}$	$\frac{1}{1}$	Cyperus sp.					************			0.00	
salt grass	FACW	1	1	1	Distichlis spicata								0.00	
spikeweed	OBL	1	1	1	Eleocharis sp.					***************************************			0.00	
common waterweed hairy willowherb	OBL FACW	1	1	1	Elodea canadensis Epilobium ciliatum								0.00	
seaside heliotrope	OBL	1	1	1	Heliotropium curassavicum								0.00	
water penny	OBL	1	1	1	Hydrocotyl sp.								0.00	
Baltic rush	OBL	1	1	1	Juncus balticus		1.5			2.5	1.5	97.5	0.00	
soft rush iris-leaved rush	OBL OBL	1	1	$\frac{1}{1}$	Juncus effusus Juncus xiphioides		15			2.5	15	87.5	17.14 0.00	
creeping wildrye	FAC+	1	1	1	Leymus triticoides						2.5		0.36	
water primrose	OBL	1	1	1	Ludwigia peploides								0.00	
swamp smartweed Fremont cottonwood	OBL FACW	1	1	$\frac{1}{1}$	Polygonum hydropiperoides Populus fremontii	37.5	62.5		62.5	15	15		0.00 27.50	
California rose	FAC+	1	1	1	Rosa californica	37.3	02.3		02.3	13	13		0.00	
gray willow	OBL	1	1	1	Salix exigua								0.00	
Gooding willow	OBL	1	1	1	Salix goodingii								0.00	
red willow arroyo willow	FACW+	1	$\frac{1}{1}$	$\frac{1}{1}$	Salix laevigata Salix lasiolepis								0.00	
tule	OBL	1	1	1 1	Scirpus acutus								0.00	
three square	OBL	1	1	1	Scirpus americanus								0.00	
salt marsh bullrush	OBL	1	1	1	Scirpus robustus								0.00	
southern cattail broad-leaved cattail	OBL OBL	1	1	1 1	Typha domingensis Typha latifolia			2.5					0.00 0.36	
stinging nettle	FACW	1	1	1	Urtica dioica ssp holosericea			4.3					0.00	
fat hen	FACW	1	1	0	Atriplex triangularis								0.00	
horseweed	FAC	1	1	0	Conyza canadensis								0.00	
toad rush broad-leaved pepperweed	FACW+	1	0	0	Juncus bufonius Lepidium latifolium		***************************************			200000000000000000000000000000000000000		000000000000000000000000000000000000000	0.00	
bird's-foot trefoil	FACW	1	0	1	Lepiaium iaujoiium Lotus corniculatus	37.5							5.36	
dallisgrass	FAC	1	0	1	Paspalum dilatatum			2.5					0.36	
Vasey grass	OBL	1	0	1	Paspalum urvillei								0.00	
Kentucky bluegrass dooryard knotweed	FAC FAC	1	0	$\frac{1}{1}$	Poa pratensis Polygonum arenastrum		0.1						0.00	
water beard grass	FACW	1	0	$\frac{1}{1}$	Polypogon viridis		0.1						0.00	
green dock	FACW	1	0	1	Rumex conglomeratus								0.00	
curly dock	FACW-	1	0	1	Rumex crispus						0.1		0.01	
Fiddleleaf dock Himalaya blackberry	FAC FACW	1	0	1	Rumex pulcher Rubus discolor								0.00	
Pacific bent grass	FACW*	1	0	0	Agrostis avenacea					***************************************		000000000000000000000000000000000000000	0.00	
Slender centaury	FACW	1	0	0	Centaurium tenuiflorum								0.00	
lamb's quarters	FAC	1	0	0	Chenopodium album								0.00	
poison hemlock water hyacinth	FACW OBL	1	0	0	Conium maculatum Eichhornia crassipes								0.00	
weedy cudweed	FACW-	1	0	0	Gnaphalium luteo-album								0.00	
Seaside barley	FAC	1	0	0	Hordeum marinum								0.00	
prickly lettuce	FAC OBL	1	0	0	Lactuca serriola								0.00	
bearded sprangletop Italian ryegrass	FAC	1	0	0	Leptochloa fiscicularis Lolium multiflorum								0.00	
hyssop loosestrife	FACW	1	0	0	Lythrum hyssopifolia								0.00	
India sweet-clover	FAC	1	0	0	Melilotus indica		2.5		0.1				0.37	
bristly ox-tongue	FAC FACW-	1	0	0	Picris echioides					0.1			0.01 0.00	
annual blue-grass rabbitfoot grass	FACW+	1	0	0	Poa annua Polypogon monspeliensis	2.5	37.5					15	7.86	
coyote bush	UPL	0	1	1	Baccharis pilularis								0.00	
big leaf mistletoe	UPL	0	1	1	Phoradenron macrophyllum	000000000000000000000000000000000000000	000000000000000000000000000000000000000	000000000000000000000000000000000000000		***************************************	000000000000000000000000000000000000000	100000000000000000000000000000000000000	0.00	
parched willowherb	UPL UPL	0	1	0	Epilobium brachycarpum		2.5		2.5	0.1			0.37	
Bindweed Bermuda grass	FACU	0	$\frac{0}{0}$	$\frac{1}{1}$	Convolvulus arensis Cynoda dactylis				2.5				0.36 0.00	
wild oat	UPL	0	0	0	Avena fatua	0.1	000000000000000000000000000000000000000		0.1	15		300000000000000000000000000000000000000	2.17	
ripgut grass	UPL	0	0	0	Bromus diandrus		2.5		2.5				0.71	
soft chess Spanish Brome	FACU- UPL	0	0	0	Bromus hordeaceus Bromus madritensis	15	15 2.5		37.5	37.5		15	17.14 0.36	
Spanish Brome Italian thistle	UPL	0	0	0	Carduus pycnocephalus		2.3			0.1	2.5		0.36	
Big heron bill	FACU	0	0	0	Erodium botrys		2.5						0.36	
Rattail sixweeks grass	UPL	0	0	0	Festuca myuros								0.00	
Yellow sneezeweed summer mustard	FACU UPL	0	0	0	Helenium amarum Hirschfeldia incana								0.00	
Foxtail barley	FACU	0	0	0	Hirscnfeldia incana Hordeum murinum								0.00	
mallow	UPL	0	0	0	Malva sp								0.00	
pineapple weed	FACU	0	0	0	Matricaria matricarioides								0.00	
Bur clover white sweet-clover	FACU+	0	0	0	Medicago polymorpha Melilotus alba								0.00	
White sweet-clover Wild radish	UPL	0	0	0	Raphanus sativus								0.00	
common groundsel	NI	0	0	0	Senecio vulgaris								0.00	
common sow-thistle	NI	0	0	0	Sonchus oleraceus								0.00	
				<u> </u>	(Sp. Rich.) Number of species:	6	12	4	9	9	6	4	7.14	
# # # 4000:	mid-point	S			Number of native species:	2	4	3	4	4	4	2	3.29	
5=75-100%	5==87.5			 	Total veg cover	130.1	195.1	92.6	180.3	122.8	122.6	132.5	139.43	
4=50-75%	4==62.5				elative veg cover (=100-unveg):	100.0	85.0	100.0	100.0	100.0	100.0	99.9	97.8	
3=25-50%	3==37.5			<u> </u>	∑ cover of all hydrophytic spp:	115.0	170.1	92.6	137.7	70.1	120.1	117.5	117.6	
2=5-25%	2==15			<u> </u>	\sum cover of all native spp:	75.0	132.5	90.1	137.6	70.1	120.00	102.5	104.0	
1=many-5%	1==2.5				\sum cover of all perennial spp:	112.5	130.1	92.6	140.1	70.0	120.1	102.5	109.7	
d=few	d==0.1			<u> </u>	relative cover of hydrophytes:	88%	87%	100%	76%	57%	98%	89%	85%	
r=one	r==0.01			<u> </u>	relative cover of natives:	58%	68%	97%	76%	57%	98%	77%	76%	
					relative cover of perennials:	86%	67%	100%	78%	57%	98%	77%	80%	

		T T 1	1		1	8			hin Vaal):	II. land		
		Hydr	ophyte Nativ		Plot		5	9	hin Kee l	Aiparian 21	∪piand 28	35	Ave.
Common Name	ROIND		1 vaca v	Per	Unveg	<u> </u>			15	21	20	33	2.14
yerba mansa	OBL	1	1	1	Anemopsis californica								0.00
slender aster Santa Barbara sedge	FACW FACW	1	1	1 1	Aster subulatus var. ligulatus Carex barbarae								0.00
clustered field sedge	FACW-	1	$\frac{1}{1}$	$\frac{1}{1}$	Carex praegracilis								0.00
brown dogwood	FACW	1	1	1	Cornus glabrata								0.00
tall flat-sedge	FACW	1	1	1	Cyperus eragrostis								0.00
Sedge sp.	FACW FACW	1	1 1	1	Cyperus sp. Distichlis spicata								0.00
salt grass spikeweed	OBL	1	1	1	Eleocharis spicaia								0.00
common waterweed	OBL	1	$\frac{1}{1}$	$\frac{1}{1}$	Elodea canadensis								0.00
hairy willowherb	FACW	1	1	1	Epilobium ciliatum								0.00
seaside heliotrope	OBL	1	1	1	Heliotropium curassavicum								0.00
water penny Baltic rush	OBL OBL	1	1	$\frac{1}{1}$	Hydrocotyl sp. Juncus balticus	-		ļ	ļ				0.00
soft rush	OBL	1	1	1	Juncus effusus				2.5				0.00
iris-leaved rush	OBL	1	1 1	$\frac{1}{1}$	Juncus xiphioides	 			2.3				0.00
creeping wildrye	FAC+	1	1	1	Leymus triticoides	87.5	87.5	87.5	37.5	87.5	87.5	87.5	80.36
water primrose	OBL	1	1	1	Ludwigia peploides								0.00
swamp smartweed	OBL	1	1	1	Polygonum hydropiperoides				0.1				0.00
Fremont cottonwood California rose	FACW FAC+	1	1	1	Populus fremontii Rosa californica				0.1				0.01
gray willow	OBL	1	+ 1	$\frac{1}{1}$	Salix exigua								0.00
Gooding willow	OBL	1	1	1	Salix goodingii								0.00
red willow	FACW+	1	1	1	Salix laevigata								0.00
arroyo willow	FACW	1	1	1	Salix lasiolepis								0.00
tule	OBL	1	1	1	Scirpus acutus	<u> </u>		<u> </u>					0.00
three square salt marsh bullrush	OBL OBL	1	1 1	1 1	Scirpus americanus Scirpus robustus	-						 	0.00
southern cattail	OBL	1	1	1	Typha domingensis	 			-				0.00
broad-leaved cattail	OBL	1	1	1	Typha latifolia	<u> </u>							0.00
stinging nettle	FACW	1	1	1	Urtica dioica ssp holosericea								0.00
fat hen	FACW	1	1	0	Atriplex triangularis								0.00
horseweed tood rush	FAC FACW+	1	$\frac{1}{1}$	0	Conyza canadensis	-	 	 	 			ļl	0.00
toad rush broad-leaved pepperweed	FACW+	1	0	1	Juncus bufonius Lepidium latifolium								0.00
bird's-foot trefoil	FAC	1	0	1	Lotus corniculatus		2.5						0.36
dallisgrass	FAC	1	0	1	Paspalum dilatatum			 					0.00
Vasey grass	OBL	1	0	1	Paspalum urvillei								0.00
Kentucky bluegrass	FAC	1	0	1	Poa pratensis	ļ							0.00
dooryard knotweed water beard grass	FAC FACW	1	0	1 1	Polygonum arenastrum Polypogon viridis								0.00
green dock	FACW	1	0	1	Rumex conglomeratus								0.00
curly dock	FACW-	1	10	$\frac{1}{1}$	Rumex crispus	-							0.00
Fiddleleaf dock	FAC	1	0	1	Rumex pulcher								
Himalaya blackberry	FACW	1	0	1	Rubus discolor								0.00
Pacific bent grass	FACW*	1	0	$\frac{1}{0}$	Agrostis avenacea	ļ							0.00
Slender centaury lamb's quarters	FAC	1 1	0	$\frac{1}{0}$	Centaurium tenuiflorum Chenopodium album								0.00
poison hemlock	FACW	1	0	$\frac{1}{0}$	Conium maculatum								0.00
water hyacinth	OBL	1	0	0	Eichhornia crassipes								0.00
weedy cudweed	FACW-	1	0	0	Gnaphalium luteo-album								0.00
Seaside barley	FAC	1	$\frac{1}{0}$	$\frac{1}{0}$	Hordeum marinum	ļ		ļ					0.00
prickly lettuce bearded sprangletop	FAC OBL	1	0	0	Lactuca serriola Leptochloa fiscicularis								0.00
Italian ryegrass	FAC	1	0	1 0	Lolium multiflorum								0.00
hyssop loosestrife	FACW	1	0	0	Lythrum hyssopifolia			<u> </u>					0.00
India sweet-clover	FAC	1	0	0	Melilotus indica		15						2.14
bristly ox-tongue	FAC	1	0	0	Picris echioides								0.00
annual blue-grass rabbitfoot grass	FACW- FACW+	1	0	0	Poa annua Polypogon monspeliensis								0.00
covote bush	UPL	0	1	$\frac{1}{1}$	Baccharis pilularis						000000000000000000000000000000000000000		0.00
big leaf mistletoe	UPL	0	1	$\frac{1}{1}$	Phoradenron macrophyllum			<u> </u>					0.00
parched willowherb	UPL	0	1	0	Epilobium brachycarpum	000000000000000000000000000000000000000				2.5	0.1		0.37
Bindweed	UPL	0	0	1	Convolvulus arvensis						0.1		0.01
Bermuda grass wild oats	FACU UPL	0	0	1 0	Cynoda dactylis		ļ		2.5 0.1				0.36 0.01
vild oats ripgut grass	UPL	0	0	0	Avena fatua Bromus diandrus			 	0.1				0.01
soft chess	FACU-	0	0	0	Bromus hordeaceus	37.5	37.5	37.5	†	37.5	37.5	0.1	26.80
Spanish Brome	UPL	0	0	0	Bromus madritensis								0.00
Italian thistle	UPL	0	0	0	Carduus pycnocephalus	ļ		0.1			0.1		0.03
Big heron bill Rattail sixweeks grass	FACU UPL	0	0	$\frac{1}{0}$	Erodium botrys	-							0.00
Yellow sneezeweed	FACU	0	$\frac{0}{0}$	$\frac{0}{0}$	Festuca myuros Helenium amarum	 		 					0.00
summer mustard	UPL	0	0	0	Hirschfeldia incana	†	 	†	†				0.00
Foxtail barley	FACU	0	0	0	Hordeum murinum								0.00
mallow	UPL	0	0	0	Malva sp								0.00
pineapple weed	FACU	0	0	0	Matricaria matricarioides								0.00
Bur clover white sweet-clover	FACU+	0	0	0	Medicago polymorpha Melilotus alba	 			 				0.00
Wild radish	UPL	0	0	0	Raphanus sativus	 		 				 	0.00
common groundsel	NI	0	0	0	Senecio vulgaris								0.00
common sow-thistle	NI	0	0	0	Sonchus oleraceus								0.00
					(Sp. Rich.) Number of species:	2.00	4.00	3.00	5.00	3.00	5.00	2.00	3.43
	mid-points	S			Number of native species:	1.00	1.00	1.00	3.00	2.00	2.00	1.00	1.57
5=75-100%	5==87.5				Total veg cover	125	142.5	125.1	42.7	127.5	125.3	87.6	110.81
4=50-75%	4==62.5			r	elative veg cover (=100-unveg):	100.0	100.0	100.0	85.0	100.0	100.0	100.0	97.86
3=25-50%	3==37.5			8	∑ cover of all hydrophytic spp:	87.5	105.0	87.5	40.1	87.5	87.5	87.5	83.23
2=5-25%	2==15				\sum cover of all native spp:	87.5	87.5	87.5	40.1	90.0	87.6	87.5	81.10
1=many-5%	1==2.5				\sum cover of all perennial spp:	87.5	90.0	87.5	42.6	87.5	87.6	87.5	81.46
d=few	d=0.1				relative cover of hydrophytes:	70%	74%	70%	94%	69%	70%	100%	78%
r=one	r==0.01			—	relative cover of natives:	70%	61%	70%	94%	71%	70%	100%	77%
1-one	10.01			 		/ 070	 	 	 	 	}	10070	
				1	relative cover of perennials:	70%	63%	70%	100%	69%	70%	100%	77%